

S'ESSA NON PASSA PER LE MATEMATICHE DIMOSTRAZIONI
LEONARDO DA VINCI

vol. 4

no. 3-4

2016

MATHEMATICS AND MECHANICS *of* Complex Systems

GEOFFREY R. GRIMMETT

THE WORK OF LUCIO RUSSO ON PERCOLATION



THE WORK OF LUCIO RUSSO ON PERCOLATION

GEOFFREY R. GRIMMETT

Dedicated in friendship to Lucio Russo

The contributions of Lucio Russo to the mathematics of percolation and disordered systems are outlined. The context of his work is explained, and its ongoing impact on current work is described and amplified.

1. A personal appreciation

Prior to his mid-career move to the history of science in the early 1990s, Lucio Russo enjoyed a very successful and influential career in the theory of probability and disordered systems, in particular of percolation and the Ising model. His ideas have shaped these significant fields of science, and his name will always be associated with a number of fundamental techniques of enduring importance.

The author of this memoir is proud to have known Lucio in those days, and to have profited from his work, ideas, and company. He hopes that this brief account of some of Lucio's results will stand as testament to the beauty and impact of his ideas.

2. Scientific summary

Lucio Russo has worked principally on the mathematics of percolation, that is, of the existence (or not) of infinite connected clusters within a disordered spatial network. The principal model in this field is the so called *percolation model*, introduced to mathematicians by Broadbent and Hammersley in 1957, [14]. Consider, for definiteness, the hypercubic lattice \mathbb{Z}^d with $d \geq 2$, and let $p \in [0, 1]$. We declare each edge to be *open* with probability p and *closed* otherwise, and different edges receive independent states. The main questions are centred around the existence (or not) of an infinite open component in \mathbb{Z}^d . It turns out that there exists a critical probability $p_c = p_c(d)$ such that no infinite open cluster exists when $p < p_c$, and there exists a unique such cluster when $p > p_c$. (It is not still known which of these two occurs when $p = p_c$ for general d , specifically when $3 \leq d \leq 10$. See [20].)

Communicated by Raffaele Esposito.

MSC2010: 60K35, 82B20.

Keywords: percolation, Ising model, Russo's formula, RSW inequality, box crossing, approximate zero-one law, influence, sharp threshold.

Let C be the open cluster of \mathbb{Z}^d containing the origin. Two functions that play important roles in the theory are the *percolation probability* θ and the *mean cluster size* χ given by

$$\theta(p) = P_p(|C| = \infty), \quad \chi(p) = E_p|C|,$$

where P_p and E_p are the appropriate product measure and expectation. The above model is the *bond* percolation model; the *site* percolation model is defined similarly, with sites being open/closed. A fairly recent account of percolation may be found in [25].

The question was raised in 1960 (by Harris, [31]) of whether or not $p_c(2) = \frac{1}{2}$, and the search for a rigorous proof attracted a number of fine mathematicians into the field, including Lucio. Several important partial results were proved, culminating in 1980 with Kesten’s complete proof that $p_c(2) = \frac{1}{2}$, [37]. The interest of the community then migrated towards the case $d \geq 3$, before returning firmly to $d = 2$ with the 2001 proof by Smirnov, [44; 45], of Cardy’s formula.

Lucio contributed a number of fundamental techniques to percolation theory during the period 1978–1988, and the main purpose of the current paper is to describe these and to explore their significance. We mention Russo’s formula, the Russo–Seymour–Welsh (RSW) inequalities, his study of percolation surfaces in three dimensions, and of the uniqueness of the infinite open cluster, and finally Russo’s approximate zero–one law. Russo’s formula and RSW theory have proved of especially lasting value in, for example, recent developments concerning conformal invariance for critical percolation.

In Section 8, we mention some of Lucio’s results concerning percolation of $+/-$ spins in the two-dimensional Ising model. It was quite a novelty in the 1970s to use percolation as a tool to understand long-range order in the Ising model. Indeed, Lucio’s work on the percolation model was motivated in part by his search for rigorous results in statistical mechanics. His approach to the Ising model has been valuable in two dimensions. In more general situations, the correct geometrical model has been recognised since to be the *random-cluster model* of Fortuin and Kasteleyn (see [26]).

This short account is confined to Lucio’s contributions to percolation, and does not touch on his work lying closer to ergodic theory and dynamical systems, namely [R2; R6; R8; R11], and neither does it refer to the paper [R7]. A comprehensive list of Lucio’s mathematical publications, taken from MathSciNet, may be found at the end of this paper.

Results from Lucio’s work will be described here using ‘modern’ notation. No serious attempt is made to include comprehensive citations of the related work of others.

3. Russo's formula

Let $\Omega = \{0, 1\}^E$ where E is finite, and let P_p be product measure on the partially ordered set Ω with density $p \in [0, 1]$. An event $A \subseteq \Omega$ is called *increasing* if:

$$\omega \in A, \omega \leq \omega' \Rightarrow \omega' \in A.$$

Let $\omega \in \Omega$. An element $e \in E$ is called *pivotal* for an increasing event A if $\omega_e \notin A$ and $\omega^e \in A$, where ω^e and ω_e are obtained from ω by varying the state of the edge e thus:

$$\omega_e(f) = \begin{cases} 0 & \text{if } f = e, \\ \omega(f) & \text{if } f \neq e, \end{cases} \quad \omega^e(f) = \begin{cases} 1 & \text{if } f = e, \\ \omega(f) & \text{if } f \neq e. \end{cases}$$

In other words, e is said to be pivotal for A if the occurrence of A depends on the state of e .

Theorem 3.1 (Russo's formula, [R14]). *Let A be an increasing event. We have*

$$\frac{d}{dp} P_p(A) = \sum_{e \in E} P_p(e \text{ is pivotal for } A).$$

Similar techniques are encountered independently in related fields. For example, Russo's formula is essentially equation (4.4) of Barlow and Proschan's book [7, p. 212] on reliability theory. Such a formula appeared also in the work of Margulis, [39], in the Russian literature. A characteristic of Lucio's work is the geometric context of the formula when applied in situations such as percolation, and it is in this context that Lucio's name is prominent. In a typical application to percolation, one uses the geometrical characteristics of the event $\{e \text{ is pivotal for } A\}$ to derive differential inequalities for $P_p(A)$.

Russo's formula is key to the study of geometrical probability governed by a product measure. It has so many applications that it is a challenge to single out any one. We mention here its use in the derivation of exact values for critical exponents in two dimensions, [38; 46].

Similarly, extensions of Russo's formula have been central in several related fields, including but not limited to the contact model [10, Thm 2.13], continuum percolation [21; 34], and the random-cluster model [11, Prop. 4].

4. Russo–Seymour–Welsh inequalities

For twenty years from about 1960 to 1980, mathematicians attempted to prove that the critical probability p_c of bond percolation on the square lattice satisfies $p_c = \frac{1}{2}$. This prominent open problem was in the spirit of that of the critical temperature of the Ising model, resolved in 1944 by Onsager, [42]. Harris [31] showed how to use duality to obtain $p_c \geq \frac{1}{2}$, but the corresponding upper bound was elusive. Then, in

1978, a powerful technique emerged in independent and contemporaneous work of Lucio, [R12], and Seymour and Welsh, [43]. It has come to be known simply as ‘RSW’.

Consider bond percolation with density p on the square lattice \mathbb{Z}^2 . A *left–right crossing* of a rectangle B is an open path in B which joins some vertex on its left side to some vertex on its right side. For positive integers m and n , we define the rectangle

$$B(m, n) = [0, 2m] \times [0, 2n],$$

and let $\text{LR}(m, n)$ be the event that there exists a left–right crossing of $B(m, n)$.

Lemma 4.1 (Russo–Seymour–Welsh (RSW), [R12; 43]). *Let $p \in (0, 1)$. We have*

$$P_p(\text{LR}(\tfrac{3}{2}n, n)) \geq (1 - \sqrt{1 - \tau})^3,$$

where $\tau = P_p(\text{LR}(n, n))$.

This fundamental but superficially innocuous lemma implies that, if the chance of crossing a square is bounded from 0 uniformly in its size, then so is the chance of crossing a rectangle with aspect ratio $\frac{3}{2}$. Using the self-duality of \mathbb{Z}^2 , we have as input to the RSW lemma that

$$P_{\frac{1}{2}}(\text{LR}(n, n)) \geq \tfrac{1}{2}. \quad (4.1)$$

Let A_n be the event that the annulus $[-3n, 3n]^2 \setminus [-n, n]^2$ contains an open cycle with the origin in the bounded component of its complement in \mathbb{R}^2 . Using elementary geometrical arguments and the FKG inequality, it follows by the RSW lemma and (4.1) that there exists $\sigma > 0$ such that $P_p(A_n) \geq \sigma$ for $n \geq 1$ and $p \geq \frac{1}{2}$.

The RSW lemma and the ensuing annulus inequality have proved to be key to the study of percolation in two dimensions. In common with other useful methods of mathematics, there is now a cluster of related inequalities, see for example [12], [27, Sect. 5.5], and [50, Chap. 5].

RSW methods were used by their discoverers to make useful but incomplete progress towards proving that $p_c = \frac{1}{2}$, and they played a role in Kesten’s full proof, [37]. (The principle novelty of Kesten’s paper was a bespoke theory of sharp threshold, see Section 5.) More precisely, they led to the following result, which is presented in terms of *site* percolation on the square lattice \mathbb{Z}^2 and its *matching lattice* \mathbb{Z}_*^2 , derived by adding the two diagonals to each face of \mathbb{Z}^2 .

Theorem 4.2 (Russo, [R12]). *Consider site percolation on the square lattice \mathbb{Z}^2 . The critical points*

$$p_c = \sup\{p : \theta(p) = 0\}, \quad \pi_c = \sup\{p : \chi(p) < \infty\},$$

satisfy

$$p_c + \pi_c^* = 1, \quad p_c^* + \pi_c = 1, \quad (4.2)$$

where an asterisk denotes the corresponding values on the matching lattice.

The parallel work of Seymour and Welsh, [43], was directed at the bond model on \mathbb{Z}^2 , of which the dual model lies on a translate of \mathbb{Z}^2 . Following Kesten's proof of $p_c(\mathbb{Z}^2) = \frac{1}{2}$ for bond percolation, Lucio revisited Theorem 4.2 in [R14] with a proof that $\pi_c^* = p_c^*$, and the consequent improvement of (4.2), namely $p_c + p_c^* = 1$. He also completed the proof, begun in [R12], that θ (and, similarly, the dual percolation probability θ^*) is a continuous function on $[0, 1]$. Continuity in two dimensions has since been extended to general percolation models (see, for example, [25, Sect. 8.3]).

RSW theory is now recognised as fundamental to rigorous proofs of conformal invariance of critical two-dimensional percolation and all that comes with that. The proof of Cardy's formula, [44; 45], provides a major illustration. It was observed by Aizenman and Burchard, [3], that certain connection probabilities belong to a space of uniformly Hölder functions. Since this space is compact, such functions have subsequential limits as the mesh of the lattice approaches 0. The above Hölder property is proved using annulus inequalities.

Indeed the power of RSW arguments extends beyond percolation to a host of problems involving two-dimensional stochastic geometry, such as the FK-Ising model [18] and Voronoi percolation [49]. In addition, RSW theory provides one of the main techniques for the proof by Beffara and Duminil-Copin, [9], that the random-cluster model on \mathbb{Z}^2 with cluster-weighting parameter $q \geq 1$ has critical value $p_c(q) = \sqrt{q}/(1 + \sqrt{q})$. We retrieve Kesten's theorem by setting $q = 1$.

5. Approximate zero–one law

Kolmogorov's zero–one law may be stated as follows. Consider the infinite product space $\Omega = \{0, 1\}^{\mathbb{N}}$ endowed with the product σ -algebra and the product measure P_p . If A is an event that is independent of any finite subcollection $\{\omega(e) : e \in E\}$, $E \subseteq \mathbb{N}$, $|E| < \infty$, then $P_p(A)$ equals either 0 or 1. It follows that, for an increasing event A , there exists $p_0 \in [0, 1]$ such that

$$P_p(A) = \begin{cases} 0 & \text{if } p < p_0, \\ 1 & \text{if } p > p_0. \end{cases}$$

This law is intrinsically an *infinite-volume* effect, in that the index set is the infinite set \mathbb{N} . Lucio posed the farsighted question in [R15] of whether there exists a *finite volume* version of this result, and this led him to his 'approximate zero–one law', following.

Let $\Omega = \{0, 1\}^E$ where E is finite, and let P_p be product measure on the partially ordered set Ω with density $p \in [0, 1]$. The *influence* $I_{A,p}(e)$ of $e \in E$ on the event $A \subseteq \Omega$ is defined by

$$I_{A,p}(e) = P_p(1_A(\omega_e) \neq 1_A(\omega^e)),$$

where 1_A denotes the indicator function of A . When A is increasing, this may be written

$$I_{A,p}(e) = P_p(\omega_e \notin A, \omega^e \in A) = P_p(e \text{ is pivotal for } A). \quad (5.1)$$

Theorem 5.1 (Russo's approximate zero-one law, [R15]). *For $\epsilon > 0$, there exists $\eta > 0$ such that, if A is an increasing event and*

$$I_{A,p}(e) < \eta, \quad e \in E, \quad p \in [0, 1], \quad (5.2)$$

then there exists $p_0 \in [0, 1]$ such that

$$P_p(A) \begin{cases} \leq \epsilon & \text{if } p < p_0 - \epsilon, \\ \geq 1 - \epsilon & \text{if } p > p_0 + \epsilon. \end{cases} \quad (5.3)$$

This result was motivated by a desire to generalise certain results for box-crossing probabilities in percolation. Its impact extends far beyond percolation, and it is a precursor of a more recent theory, pioneered by Kahn, Kalai, Linial, [35] and Talagrand, [47; 48], of influence and sharp threshold. It is proved at [47, Thm 1.1] that there exists an absolute constant $c > 0$ such that, for $p \in (0, 1)$ and an increasing event A ,

$$\sum_{e \in E} I_{A,p}(e) \geq \left(\frac{c}{p(1-p) \log[2/(p(1-p))]} \right) P_p(A)(1 - P_p(A)) \log(1/m_p), \quad (5.4)$$

where

$$m_p = \max\{I_{A,p}(e) : e \in E\}.$$

It follows that, when $p \in (0, 1)$ and A is increasing,

$$\sum_{e \in E} I_{A,p}(e) \geq c' P_p(A)(1 - P_p(A)) \log(1/m_p), \quad (5.5)$$

where $c' > 0$ is an absolute constant.

Amongst the implications of (5.5) is a quantification of the relationship between ϵ and η in Theorem 5.1. Suppose (5.2) holds with $\eta \in (0, 1)$, so that $m_p \leq \eta$. By (5.5) and Russo's formula,

$$\frac{d}{dp} P_p(A) \geq c' P_p(A)(1 - P_p(A)) \log(1/\eta). \quad (5.6)$$

Choose p_0 such that $P_{p_0}(A) = \frac{1}{2}$, and integrate (5.6) to obtain

$$P_p(A) \begin{cases} \leq \epsilon_p & \text{if } p < p_0, \\ \geq 1 - \epsilon_p & \text{if } p > p_0, \end{cases} \quad (5.7)$$

with

$$\epsilon_p = \frac{1}{1 + (1/\eta)^{c'|p-p_0|}}.$$

Such inequalities have found numerous applications in percolation and related topics, see for example [12; 19; 24]. They have been extended to general product measures, [13; 28], and to probability measures satisfying the FKG lattice condition, [23]. Recent overviews include [27, Chap. 4] and [36].

6. Percolation in dimension $d \geq 3$

In 1983, Lucio spent a sabbatical at Princeton University. His work during that period led to two significant publications [R1; R3] on aspects of percolation in three dimensions. The first of these caused quite a stir in the community at the time of its appearance, largely since most work until then had been for models in only two dimensions. Whereas the dual of a bond model in two dimensions is another bond model, the dual model in three or more dimensions is a ‘plaquette’ model. Since the topology of surfaces of plaquettes is much more complicated than that of paths, the ensuing percolation duality poses a number of challenging topological questions.

The authors of [R1] consider bond percolation on \mathbb{Z}^3 with density p , together with its dual ‘plaquette’ model on $\mathbb{Z}_*^3 := \mathbb{Z}^3 + (\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$. A *plaquette* is a unit square with vertices in \mathbb{Z}_*^3 , and its bounding lines are *edges* of \mathbb{Z}_*^3 . Each edge e of \mathbb{Z}^3 intersects a unique plaquette Π_e , and Π_e is termed *occupied* if and only if e is closed (and *unoccupied* otherwise). Thus, a plaquette is occupied with probability $1 - p$. For any collection F of plaquettes, the boundary ∂F is defined to be set of edges of \mathbb{Z}_*^3 belonging to an *odd* number of members of F .

Let γ be a cycle of \mathbb{Z}_*^3 . The main results of [R1] concern the probability there exists a set F of occupied plaquettes which spans γ in the sense that $\partial F = \gamma$. For simplicity, we shall suppose here that γ is a $m \times n$ rectangle of the x/y plane, and we denote the above event as W_γ . Note that γ has area mn and perimeter $2(m+n)$.

Theorem 6.1 (Aizenman, Chayes, Chayes, Fröhlich, Russo, [R1]). *There exist constants $\pi_c, \rho_c \in (0, 1)$ such that*

$$-\log P_p(W_\gamma) \sim \begin{cases} \alpha mn & \text{if } 1 - p < \pi_c, \\ \beta(m+n) & \text{if } 1 - p > \rho_c, \end{cases}$$

where $\alpha, \beta > 0$ depend on p , and the asymptotic relation is as $m, n \rightarrow \infty$.

The constants π_c , ρ_c are the critical densities of the bond percolation model on \mathbb{Z}^3 given by

$$\pi_c = \sup\{p : \chi(p) < \infty\}, \quad \rho_c = \lim_{k \rightarrow \infty} \widehat{p}_c(k),$$

where $\widehat{p}_c(k)$ is the *slab critical point*

$$\widehat{p}_c(k) = \sup\{p : P_p(0 \leftrightarrow \infty \text{ in } [0, \infty)^2 \times [0, k]) = 0\}.$$

It was conjectured in [R1] that $\pi_c = p_c = \rho_c$. The first equality was proved later in [2; 41], and the second in [8; 30].

There are only few percolation models on finite-dimensional lattices for which the numerical values of the critical probabilities are known exactly, and all such exact results are in two dimensions only (see, for example, [29]). In contrast, quite a lot of work has been devoted to obtaining rigorous upper and lower bounds for critical probabilities, and there is a host of numerical estimates.

Consider site percolation on the simple cubic lattice \mathbb{Z}^3 . By a comparison with the site model on the triangular lattice, Lucio has shown (with Campanino, in [R3]) that $p_c \leq \frac{1}{2}$. (See also [40].) They obtained also the strict inequality, with a distinctly more complicated argument.

Theorem 6.2 (Campanino, Russo, [R3]). *The critical probability of site percolation on \mathbb{Z}^3 satisfies $p_c < \frac{1}{2}$.*

The point of this work was to show that, in a neighbourhood of $p = \frac{1}{2}$, there is coexistence of infinite open and infinite closed clusters in \mathbb{Z}^3 . The corresponding statement for $d = 2$ is, of course, false, in that coexistence occurs for no value of p .

Theorem 6.2 may still be the best rigorous upper bound that is currently known for p_c . By examining its proof, one may calculate a small $\epsilon > 0$ such that $p_c < \frac{1}{2} - \epsilon$. It is expected that $p_c \approx 0.31$.

7. Uniqueness of the infinite open cluster

Let I be the number of infinite open clusters of a percolation model in a finite-dimensional space. For a period in the 1980s, the ‘next’ problem was to prove that $P_p(I = 1) = 1$ in the supercritical phase (when $p > p_c$). This problem was solved by Aizenman, Kesten, and Newman [5] in 1987. Their proof seemed slightly mysterious at the time, and it was simplified by Lucio in the jointly written paper [R9]. The key step was to show, using a large-deviation estimate present already in [5], that there is density 0 of sites that are adjacent to *two* distinct infinite clusters.

This useful argument was soon overshadowed by the beautiful proof of uniqueness by Burton and Keane, [15], of which a key step is a novel argument to show

there is density 0 of sites that are adjacent to *three* distinct infinite clusters. The proof of [15] uses translation-invariance of the underlying measure together with a property of so called ‘finite energy’, and may thus be extended to more general measures than product measures. On the other hand, since the proof uses no quantitative estimate, it yields no ‘rate’. The methods of [5; R9] provide a missing rate, and this has been useful in the later work [16; 17].

The question of uniqueness for dependent models is potentially harder, since the large-deviation estimate of [5; R9] is not available. In joint work [R10] with Gandolfi and Keane, Lucio used path-intersection arguments to show uniqueness for ergodic, positively associated measures in two dimensions, satisfying certain translation and reflection symmetries. Unlike the Burton–Keane proof, they needed no finite-energy assumption. An application of this work to quantum spin systems may be found in [6].

8. Ising model

Lucio has written three papers on the geometry of the d -dimensional Ising model, [R4; R5; R13]. In this work, he (and his coauthors) studied the relationship between properties of the infinite-volume Gibbs measures and the existence or not of an infinite cluster of either $+$ or $-$ spins (that is, of percolation in the Ising model).

The first two of these papers [R4; R5] explore a relationship between the Ising magnetization and the above percolation probability, and yield the non-existence of percolation in the high-temperature phase. This is complemented when $d = 2$ with the proof that percolation (of the corresponding spin) exists in the low-temperature phase for either of the pure infinite-volume limits μ_+ , μ_- , obtained respectively as the weak limits with $+/-$ boundary conditions. These methods were developed further in [R5], where a phase diagram was proposed for the existence of infinite clusters in the two-dimensional ferromagnetic Ising model, as a function of external field h and temperature T . The principal features of this diagram were later proved by Higuchi, [33].

One of the central problems in two dimensions of the late 1970s was to prove or disprove the statement that every infinite-volume Gibbs measure is a convex combination of the two extremal measures μ_+ , μ_- . Lucio obtained the following important result for this problem.

Theorem 8.1 (Russo, [R13]). *Any infinite-volume Gibbs measure μ , which is translation-invariant in one or both of the axial directions, is a convex combination of μ_+ and μ_- .*

Lucio proved this by considering the existence (or not) of infinite $+/-$ clusters on \mathbb{Z}^2 and its matching lattice. The full conclusion, without an assumption of partial translation-invariance, was obtained later in independent work of Aizenman,

[1], and Higuchi, [32] (see also [22]). Therefore, in two dimensions (unlike three dimensions) there exists no non-translation-invariant Gibbs measure.

More recent work on the geometrical properties of the Ising model has been centred around the random-cluster model and the random-current representation, rather than the more fundamental percolation model. See, for example, [4; 26].

Acknowledgements

This work was supported in part by the Engineering and Physical Sciences Research Council under grant EP/I03372X/1. The author thanks Alberto Gandolfi for his comments and suggestions.

Mathematical publications of Lucio Russo

- [R1] M. Aizenman, J. T. Chayes, L. Chayes, J. Fröhlich, and L. Russo, “On a sharp transition from area law to perimeter law in a system of random surfaces”, *Comm. Math. Phys.* **92**:1 (1983), 19–69.
- [R2] C. Cammarota and L. Russo, “Bernoulli and Gibbs probabilities of subgroups of $\{0, 1\}^S$ ”, *Forum Math.* **3**:4 (1991), 401–414.
- [R3] M. Campanino and L. Russo, “An upper bound on the critical percolation probability for the three-dimensional cubic lattice”, *Ann. Probab.* **13**:2 (1985), 478–491.
- [R4] A. Coniglio, C. R. Nappi, F. Peruggi, and L. Russo, “Percolation and phase transitions in the Ising model”, *Comm. Math. Phys.* **51**:3 (1976), 315–323.
- [R5] A. Coniglio, C. R. Nappi, F. Peruggi, and L. Russo, “Percolation points and critical point in the Ising model”, *J. Phys. A* **10**:2 (1977), 205–218.
- [R6] F. di Liberto, G. Gallavotti, and L. Russo, “Markov processes, Bernoulli schemes, and Ising model”, *Comm. Math. Phys.* **33** (1973), 259–282.
- [R7] G. Facchinetti and L. Russo, “A one-dimensional case of stochastic homogenization”, *Boll. Un. Mat. Ital. C* (6) **2**:1 (1983), 159–170.
- [R8] V. Franceschini and L. Russo, “Stable and unstable manifolds of the Hénon mapping”, *J. Statist. Phys.* **25**:4 (1981), 757–769.
- [R9] A. Gandolfi, G. Grimmett, and L. Russo, “On the uniqueness of the infinite cluster in the percolation model”, *Comm. Math. Phys.* **114**:4 (1988), 549–552.
- [R10] A. Gandolfi, M. Keane, and L. Russo, “On the uniqueness of the infinite occupied cluster in dependent two-dimensional site percolation”, *Ann. Probab.* **16**:3 (1988), 1147–1157.
- [R11] G. Monroy and L. Russo, “A family of codes between some Markov and Bernoulli schemes”, *Comm. Math. Phys.* **43**:2 (1975), 155–159.
- [R12] L. Russo, “A note on percolation”, *Z. Wahrsch. Verw. Gebiete* **43**:1 (1978), 39–48.
- [R13] L. Russo, “The infinite cluster method in the two-dimensional Ising model”, *Comm. Math. Phys.* **67**:3 (1979), 251–266.
- [R14] L. Russo, “On the critical percolation probabilities”, *Z. Wahrsch. Verw. Gebiete* **56**:2 (1981), 229–237.
- [R15] L. Russo, “An approximate zero-one law”, *Z. Wahrsch. Verw. Gebiete* **61**:1 (1982), 129–139.

References

- [1] M. Aizenman, “Translation invariance and instability of phase coexistence in the two-dimensional Ising system”, *Comm. Math. Phys.* **73**:1 (1980), 83–94.
- [2] M. Aizenman and D. J. Barsky, “Sharpness of the phase transition in percolation models”, *Comm. Math. Phys.* **108**:3 (1987), 489–526.
- [3] M. Aizenman and A. Burchard, “Hölder regularity and dimension bounds for random curves”, *Duke Math. J.* **99**:3 (1999), 419–453.
- [4] M. Aizenman, H. Duminil-Copin, and V. Sidoravicius, “Random currents and continuity of Ising model’s spontaneous magnetization”, *Comm. Math. Phys.* **334**:2 (2015), 719–742.
- [5] M. Aizenman, H. Kesten, and C. M. Newman, “Uniqueness of the infinite cluster and continuity of connectivity functions for short and long range percolation”, *Comm. Math. Phys.* **111**:4 (1987), 505–531.
- [6] M. Aizenman and B. Nachtergaele, “Geometric aspects of quantum spin states”, *Comm. Math. Phys.* **164**:1 (1994), 17–63.
- [7] R. E. Barlow and F. Proschan, *Mathematical theory of reliability*, Wiley, New York, 1965.
- [8] D. J. Barsky, G. R. Grimmett, and C. M. Newman, “Percolation in half-spaces: equality of critical densities and continuity of the percolation probability”, *Probab. Theory Rel. Fields* **90**:1 (1991), 111–148.
- [9] V. Beffara and H. Duminil-Copin, “The self-dual point of the two-dimensional random-cluster model is critical for $q \geq 1$ ”, *Probab. Theory Rel. Fields* **153**:3-4 (2012), 511–542.
- [10] C. E. Bezuidenhout and G. R. Grimmett, “Exponential decay for subcritical contact and percolation processes”, *Ann. Probab.* **19**:3 (1991), 984–1009.
- [11] C. E. Bezuidenhout, G. R. Grimmett, and H. Kesten, “Strict inequality for critical values of Potts models and random-cluster processes”, *Comm. Math. Phys.* **158**:1 (1993), 1–16.
- [12] B. Bollobás and O. Riordan, “The critical probability for random Voronoi percolation in the plane is $1/2$ ”, *Probab. Theory Rel. Fields* **136**:3 (2006), 417–468.
- [13] J. Bourgain, J. Kahn, G. Kalai, Y. Katznelson, and N. Linial, “The influence of variables in product spaces”, *Israel J. Math.* **77**:1-2 (1992), 55–64.
- [14] S. R. Broadbent and J. M. Hammersley, “Percolation processes, I: Crystals and mazes”, *Proc. Cambridge Philos. Soc.* **53** (1957), 629–641.
- [15] R. M. Burton and M. Keane, “Density and uniqueness in percolation”, *Comm. Math. Phys.* **121**:3 (1989), 501–505.
- [16] R. Cerf, “A lower bound on the two-arms exponent for critical percolation on the lattice”, *Ann. Probab.* **43**:5 (2015), 2458–2480.
- [17] S. Chatterjee and S. Sen, “Minimal spanning trees and Stein’s method”, 2013. [arXiv 1307.1661](https://arxiv.org/abs/1307.1661)
- [18] D. Chelkak, H. Duminil-Copin, and C. Hongler, “Crossing probabilities in topological rectangles for the critical planar FK-Ising model”, *Electron. J. Probab.* **21** (2016), 1–28.
- [19] H. Duminil-Copin and I. Manolescu, “The phase transitions of the planar random-cluster and Potts models with $q \geq 1$ are sharp”, *Probab. Theory Related Fields* **164**:3-4 (2016), 865–892.
- [20] R. Fitzner and R. van der Hofstad, “Generalized approach to the non-backtracking lace expansion”, (2015). [arXiv 1506.07969](https://arxiv.org/abs/1506.07969)
- [21] M. Franceschetti, M. D. Penrose, and T. Rosoman, “Strict inequalities of critical values in continuum percolation”, *J. Stat. Phys.* **142**:3 (2011), 460–486.

- [22] H.-O. Georgii and Y. Higuchi, “Percolation and number of phases in the two-dimensional Ising model”, *J. Math. Phys.* **41**:3 (2000), 1153–1169.
- [23] B. T. Graham and G. R. Grimmett, “Influence and sharp-threshold theorems for monotonic measures”, *Ann. Probab.* **34**:5 (2006), 1726–1745.
- [24] B. T. Graham and G. R. Grimmett, “Sharp thresholds for the random-cluster and Ising models”, *Ann. Appl. Probab.* **21**:1 (2011), 240–265.
- [25] G. R. Grimmett, *Percolation*, 2nd ed., Grundlehren der Mathematischen Wissenschaften **321**, Springer, Berlin, 1999.
- [26] G. R. Grimmett, *The random-cluster model*, Grundlehren der Mathematischen Wissenschaften **333**, Springer, Berlin, 2006.
- [27] G. R. Grimmett, *Probability on graphs*, Institute of Mathematical Statistics Textbooks **1**, Cambridge University Press, 2010.
- [28] G. R. Grimmett, S. Janson, and J. R. Norris, “Influence in product spaces”, *Adv. Appl. Probab.* **48A** (2016), 145–152.
- [29] G. R. Grimmett and I. Manolescu, “Bond percolation on isoradial graphs: criticality and universality”, *Probab. Theory Rel. Fields* **159**:1-2 (2014), 273–327.
- [30] G. R. Grimmett and J. M. Marstrand, “The supercritical phase of percolation is well behaved”, *Proc. Roy. Soc. London Ser. A* **430**:1879 (1990), 439–457.
- [31] T. E. Harris, “A lower bound for the critical probability in a certain percolation process”, *Proc. Cambridge Philos. Soc.* **56** (1960), 13–20.
- [32] Y. Higuchi, “On the absence of non-translation invariant Gibbs states for the two-dimensional Ising model”, pp. 517–534 in *Random fields* (Esztergom, Hungary, 1979), vol. I, edited by J. Fritz et al., Colloq. Math. Soc. János Bolyai **27**, North-Holland, Amsterdam, 1981.
- [33] Y. Higuchi, “A sharp transition for the two-dimensional Ising percolation”, *Probab. Theory Rel. Fields* **97**:4 (1993), 489–514.
- [34] J. Jiang, S. Zhang, and T. Guo, “Russo’s formula, uniqueness of the infinite cluster, and continuous differentiability of free energy for continuum percolation”, *J. Appl. Probab.* **48**:3 (2011), 597–610.
- [35] J. Kahn, G. Kalai, and N. Linial, “The influence of variables on Boolean functions”, pp. 68–80 in *Proceedings of the 29th annual symposium on foundations of computer science*, IEEE Computer Society, Washington, D.C., 1988.
- [36] G. Kalai and S. Safra, “Threshold phenomena and influence: perspectives from mathematics, computer science, and economics”, pp. 25–60 in *Computational complexity and statistical physics*, edited by A. G. Percus, G. Istrate, and C. Moore, Oxford University Press, 2006.
- [37] H. Kesten, “The critical probability of bond percolation on the square lattice equals $\frac{1}{2}$ ”, *Comm. Math. Phys.* **74**:1 (1980), 41–59.
- [38] H. Kesten, “Scaling relations for 2D-percolation”, *Comm. Math. Phys.* **109**:1 (1987), 109–156.
- [39] G. A. Margulis, “Probabilistic characteristics of graphs with large connectivity”, *Problemy Peredachi Informatsii* **10**:2 (1974), 101–108.
- [40] M. V. Menshikov, “Estimates for percolation thresholds for lattices in \mathbb{R}^n ”, *Dokl. Akad. Nauk SSSR* **284**:1 (1985), 36–39. In Russian; translated in *Soviet Math. Dokl.* **32**:02 (1985), 368–370.
- [41] M. V. Menshikov, “Coincidence of critical points in percolation problems”, *Dokl. Akad. Nauk SSSR* **288**:6 (1986), 1308–1311. In Russian; translated in *Soviet Math. Dokl.* **33**:03 (1986), 856–859.
- [42] L. Onsager, “Crystal statistics, I: A two-dimensional model with an order-disorder transition”, *Phys. Rev. (2)* **65** (1944), 117–149.

- [43] P. D. Seymour and D. J. A. Welsh, “Percolation probabilities on the square lattice”, pp. 227–245 in *Advances in graph theory*, vol. 3, Annals of Discrete Mathematics **3**, edited by B. Bollobás, North-Holland, Amsterdam, 1978.
- [44] S. Smirnov, “[Critical percolation in the plane: conformal invariance, Cardy’s formula, scaling limits](#)”, *C. R. Acad. Sci. Paris Sér. I Math.* **333**:3 (2001), 239–244.
- [45] S. Smirnov, “Critical percolation in the plane”, (2009). [arXiv 0909.4499](#)
- [46] S. Smirnov and W. Werner, “[Critical exponents for two-dimensional percolation](#)”, *Math. Res. Lett.* **8**:5-6 (2001), 729–744.
- [47] M. Talagrand, “[On Russo’s approximate zero-one law](#)”, *Ann. Probab.* **22**:3 (1994), 1576–1587.
- [48] M. Talagrand, “[Concentration and influences](#)”, *Israel J. Math.* **111** (1999), 275–284.
- [49] V. Tassion, “[Crossing probabilities for Voronoi percolation](#)”, *Ann. Probab.* **44**:5 (2016), 3385–3398.
- [50] W. Werner, *Percolation et modèle d’Ising*, Cours Spécialisés **16**, Société Mathématique de France, Paris, 2009.

Received 17 Apr 2016. Accepted 19 May 2016.

GEOFFREY R. GRIMMETT: g.r.grimmett@statslab.cam.ac.uk

*Statistical Laboratory, Centre for Mathematical Sciences, University of Cambridge,
Wilberforce Road, Cambridge, CB3 0WB, United Kingdom*



MATHEMATICS AND MECHANICS OF COMPLEX SYSTEMS

msp.org/memocs

EDITORIAL BOARD

ANTONIO CARCATERRA	Università di Roma "La Sapienza", Italia
ERIC A. CARLEN	Rutgers University, USA
FRANCESCO DELL'ISOLA	(CO-CHAIR) Università di Roma "La Sapienza", Italia
RAFFAELE ESPOSITO	(TREASURER) Università dell'Aquila, Italia
ALBERT FANNJIANG	University of California at Davis, USA
GILLES A. FRANCFORT	(CO-CHAIR) Université Paris-Nord, France
PIERANGELO MARCATI	Università dell'Aquila, Italy
JEAN-JACQUES MARIGO	École Polytechnique, France
PETER A. MARKOWICH	DAMTP Cambridge, UK, and University of Vienna, Austria
MARTIN OSTOJA-STARZEWSKI	(CHAIR MANAGING EDITOR) Univ. of Illinois at Urbana-Champaign, USA
PIERRE SEPPECHER	Université du Sud Toulon-Var, France
DAVID J. STEIGMANN	University of California at Berkeley, USA
PAUL STEINMANN	Universität Erlangen-Nürnberg, Germany
PIERRE M. SUQUET	LMA CNRS Marseille, France

MANAGING EDITORS

MICOL AMAR	Università di Roma "La Sapienza", Italia
CORRADO LATTANZIO	Università dell'Aquila, Italy
ANGELA MADEO	Université de Lyon-INSA (Institut National des Sciences Appliquées), France
MARTIN OSTOJA-STARZEWSKI	(CHAIR MANAGING EDITOR) Univ. of Illinois at Urbana-Champaign, USA

ADVISORY BOARD

ADNAN AKAY	Carnegie Mellon University, USA, and Bilkent University, Turkey
HOLM ALTENBACH	Otto-von-Guericke-Universität Magdeburg, Germany
MICOL AMAR	Università di Roma "La Sapienza", Italia
HARM ASKES	University of Sheffield, UK
TEODOR ATANACKOVIĆ	University of Novi Sad, Serbia
VICTOR BERDICHEVSKY	Wayne State University, USA
GUY BOUCHITTÉ	Université du Sud Toulon-Var, France
ANDREA BRAIDES	Università di Roma Tor Vergata, Italia
ROBERTO CAMASSA	University of North Carolina at Chapel Hill, USA
MAURO CARFORE	Università di Pavia, Italia
ERIC DARVE	Stanford University, USA
FELIX DARVE	Institut Polytechnique de Grenoble, France
ANNA DE MASI	Università dell'Aquila, Italia
GIANPIETRO DEL PIERO	Università di Ferrara and International Research Center MEMOCS, Italia
EMMANUELE DI BENEDETTO	Vanderbilt University, USA
BERNOLD FIEDLER	Freie Universität Berlin, Germany
IRENE M. GAMBA	University of Texas at Austin, USA
DAVID Y. GAO	Federation University and Australian National University, Australia
SERGEY GAVRILYUK	Université Aix-Marseille, France
TIMOTHY J. HEALEY	Cornell University, USA
DOMINIQUE JEULIN	École des Mines, France
ROGER E. KHAYAT	University of Western Ontario, Canada
CORRADO LATTANZIO	Università dell'Aquila, Italy
ROBERT P. LIPTON	Louisiana State University, USA
ANGELO LUONGO	Università dell'Aquila, Italia
ANGELA MADEO	Université de Lyon-INSA (Institut National des Sciences Appliquées), France
JUAN J. MANFREDI	University of Pittsburgh, USA
CARLO MARCHIORO	Università di Roma "La Sapienza", Italia
GÉRARD A. MAUGIN	Université Paris VI, France
ROBERTO NATALINI	Istituto per le Applicazioni del Calcolo "M. Picone", Italy
PATRIZIO NEFF	Universität Duisburg-Essen, Germany
ANDREY PIATNITSKI	Narvik University College, Norway, Russia
ERRICO PRESUTTI	Università di Roma Tor Vergata, Italy
MARIO PULVIRENTI	Università di Roma "La Sapienza", Italia
LUCIO RUSSO	Università di Roma "Tor Vergata", Italia
MIGUEL A. F. SANJUAN	Universidad Rey Juan Carlos, Madrid, Spain
PATRICK SELVADURAI	McGill University, Canada
ALEXANDER P. SEYRANIAN	Moscow State Lomonosov University, Russia
MIROSLAV ŠILHAVÝ	Academy of Sciences of the Czech Republic
GUIDO SWEERS	Universität zu Köln, Germany
ANTOINETTE TORDSILLAS	University of Melbourne, Australia
LEV TRUSKINOVSKY	École Polytechnique, France
JUAN J. L. VELÁZQUEZ	Bonn University, Germany
VINCENZO VESPRI	Università di Firenze, Italia
ANGELO VULPIANI	Università di Roma La Sapienza, Italia

MEMOCS (ISSN 2325-3444 electronic, 2326-7186 printed) is a journal of the International Research Center for the Mathematics and Mechanics of Complex Systems at the Università dell'Aquila, Italy.

Cover image: "Tangle" by © John Horigan; produced using the *Context Free* program (contextfreetart.org).

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing
<http://msp.org/>

© 2016 Mathematical Sciences Publishers

Special issue in honor of
Lucio Russo

Lucio Russo: A multifaceted life Raffaele Esposito and Francesco dell'Isola	197
The work of Lucio Russo on percolation Geoffrey R. Grimmett	199
"Mathematics" and "physics" in the science of harmonics Stefano Isola	213
From quantum to classical world: emergence of trajectories in a quantum system Rodolfo Figari and Alessandro Teta	235
Propagation of chaos and effective equations in kinetic theory: a brief survey Mario Pulvirenti and Sergio Simonella	255
What decides the direction of a current? Christian Maes	275
A remark on eigenvalue perturbation theory at vanishing isolation distance Fiorella Barone and Sandro Graffi	297
Some results on the asymptotic behavior of finite connection probabilities in percolation Massimo Campanino and Michele Gianfelice	311
Correlation inequalities for the Potts model Geoffrey R. Grimmett	327
Quantum mechanics: some basic techniques for some basic models, I: The models Vincenzo Grecchi	335
Quantum mechanics: some basic techniques for some basic models, II: The techniques Vincenzo Grecchi	353
On stochastic distributions and currents Vincenzo Capasso and Franco Flandoli	373
A note on Gibbs and Markov random fields with constraints and their moments Alberto Gandolfi and Pietro Lenarda	407
Quantum mechanics: light and shadows (ontological problems and epistemic solutions) Gianfausto Dell'Antonio	423
Lucio Russo: probability theory and current interests Giovanni Gallavotti	461
An attempt to let the "two cultures" meet: relationship between science and architecture in the design of Greek temples. Claudio D'Amato	471

MEMOCS is a journal of the International Research Center for
the Mathematics and Mechanics of Complex Systems
at the Università dell'Aquila, Italy.

